ELSEVIER

Physica C 263 (1996) 563–570

PHYSICA C

Author Index to Volume 263

Abe, T., see Oda, M., **263**, 241
Akazawa, S., J. Inoue and S. Maekawa, Dynamics of a carrier in the antiferromagnetic state with strong Hund coupling, **263**, 142
Akimitsu, J., M. Uehara, T. Nagata, S. Matsumoto, Y. Kitaoka, H. Takahashi and N. Môri, Metal–insulator transition in the $S=\frac{1}{2}$ spin ladder system $(Sr_{0.4}Ca_{0.6})_{14}Cu_{24-x}Co_xO_{41-\delta}$, **263**, 475
Akimitsu, J., see Nakata, H., **263**, 344
Aomine, T., see Yamasaki, Y., **263**, 317
Aomine, T., see Yasuda, T., **263**, 416
Asamitsu, A., see Tokura, Y., **263**, 544
Asaoka, H., see Ishida, T., **263**, 260
Asaoka, H., see Onodera, Y., **263**, 409
Asayama, K., see Ishida, K., **263**, 371
Asayama, K., see Magishi, K., **263**, 375
Asayama, K., see Zheng, G.-q., **263**, 367
Asayama, N., see Yamanaka, A., **263**, 516
Assaad, F.F. and M. Imada, High-frequency Hall coefficient for the two-dimensional Hubbard model, **263**, 78
Athanassopoulou, N., see Liang, W.Y., **263**, 277

Birgeneau, R.J., see Endoh, Y., **263**, 349
Braun, D.W., see Crabtree, G.W., **263**, 401

Casa, D., see Civale, L., **263**, 389
Chainani, A., see Yokoya, T., **263**, 505
Chau, R., see Maple, M.B., **263**, 490
Chi, C.C., see Tsuei, C.C., **263**, 232
Chiba, K., see Goto, T., **263**, 355
Civale, L., G. Pasquini, P. Levy, G. Nieva, D. Casa and H. Lanza, Time relaxation of persistent currents in $YBa_2Cu_3O_{7-x}$ crystals with columnar defects, **263**, 389
Cooper, J.R., see Liang, W.Y., **263**, 277
Crabtree, G.W., G.K. Leaf, H.G. Kaper, V.M. Vinokur, A.E. Koshelev, D.W. Braun, D.M. Levine, W.K. Kwok and J.A. Fendrich, Time-dependent Ginzburg–Landau simulations of vortex guidance by twin boundaries, **263**, 401

Dagotto, E., see Tsutsui, K., **263**, 107
De Andrade, M.C., see Maple, M.B., **263**, 490
Deppe, M., see Steglich, F., **263**, 498
Dessau, D.S., see Loeser, A.G., **263**, 208
Deutscher, G., see Racah, D., **263**, 218
Diver, A.J., see Mackenzie, A.P., **263**, 510
Doh, H. and S.-H.S. Salk, A Hubbard model study of antiferromagnetic ground states as a function of external magnetic field, **263**, 86
Doukan, T., see Ekino, T., **263**, 249

Ebisu, S., see Kijima, N., **263**, 523
Eder, R. and Y. Ohta, Doping dependence of the spectral function in the t–J model, **263**, 104
Eder, R., see Ohta, Y., **263**, 94
Eder, R., see Tsutsui, K., **263**, 107
Egami, M., see Yao, X., **263**, 197
Egi, T., see Gu, G.D., **263**, 180
Egi, T., see Ting, W., **263**, 185
Ekino, T., T. Doukan, H. Fujii, F. Nakamura, S. Sakita, M. Kodama and T. Fujita, Superconducting energy gap of $La_{1.85}Sr_{0.15}CuO_4$ single crystals from break-junction tunneling, **263**, 249
Emery, V.J. and S.A. Kivelson, Charge ordering in high-temperature superconductors, **263**, 44
Endoh, Y., K. Yamada, S. Wakimoto, C.H. Lee, S. Hosoya, R.J. Birgeneau, M. Greven, M.A. Kastner and G. Shirane, Spin fluctuations in superconducting $La_{1-x}Sr_xCuO_4$, **263**, 349
Enomoto, N., see Machida, K., **263**, 428
Enomoto, Y., Y. Ishikawa and S. Maekawa, Computer simulation of the magnetization process in a pulse external field, **263**, 21

Fendrich, J.A., see Crabtree, G.W., **263**, 401
Fujii, H., see Ekino, T., **263**, 249
Fujishiro, H., M. Ikebe, K. Nakasato and K. Noto, Influence of Cu site impurities on the thermal conductivity of $YBa_2Cu_3O_{7-\delta}$, **263**, 305
Fujishiro, H., see Ikebe, M., **263**, 309
Fujita, T., see Ekino, T., **263**, 249
Fujita, T., see Ikeda, S., **263**, 558
Fujita, T., see Kato, M., **263**, 482
Fujita, T., see Mackenzie, A.P., **263**, 510
Fujita, T., see Nakamura, F., **263**, 282
Fujita, T., see Yamanaka, A., **263**, 516
Fujita, T., see Yoshida, K., **263**, 519
Fujiwara, A., see Koike, Y., **263**, 329
Fujiyoshi, T., T. Sueyoshi, H. Kawamura, K. Miyahara and H. Kubota, Dynamics of flux creep in high-T_c oxide superconductors, **263**, 446
Fukami, T., see Yamasaki, Y., **263**, 317

Fukami, T., see Yasuda, T., **263**, 416
Fukase, T., M. Kamata, K. Ishizuka, T. Suzuki, T. Goto and T. Sasaki, Tilt angle effect of intrinsic flux pinning in a single crystal of $La_{1.85}Sr_{0.15}CuO_4$, **263**, 420
Fukase, T., see Goto, T., **263**, 253
Fukase, T., see Goto, T., **263**, 355
Fukase, T., see Matsuyama, A., **263**, 534
Fukase, T., see Sasaki, T., **263**, 538
Fukase, T., see Seki, K., **263**, 290
Fukuda, Y., see Nagoshi, M., **263**, 294
Fukuda, Y., see Sanada, N., **263**, 286
Fukunaga, M., see Goto, T., **263**, 355
Fukushima, K. and H. Sato, A model for high-T_c superconductivity based on charge confinement and a possible experimental verification by charge Kosterlitz–Thouless transition, **263**, 122
Fukuyama, H., Anomalous metallic state in two dimensions, **263**, 35
Fukuzumi, Y., see Uchida, S., **263**, 264
Fulde, P., see Saikawa, T., **263**, 74
Furukawa, N., Magnetic and transport properties of (La,Sr) MnO_3, **263**, 134

Gaifullin, M.B., see Matsuda, Y., **263**, 457
Gajewski, D.A., see Maple, M.B., **263**, 490
Gegenwart, P., see Steglich, F., **263**, 498
Gegenwart, P., see Takahashi, S., **263**, 30
Geibel, C., see Steglich, F., **263**, 498
Goto, T., K. Chiba, M. Fukunaga and T. Fukase, NMR study on magnetic phase transition in electron-doped cuprate $(La_{1-y}Y_y)_{2-x}Ce_xCuO_4$ with T′-type structure, **263**, 355
Goto, T., S. Nakajima, M. Kikuchi, Y. Syono and T. Fukase, Cu/Tl-NMR study on the impurity effect for Tl-based high-T_c cuprate in the over-doped region, **263**, 253
Goto, T., Critical current density of filamentary Y123 superconductor, **263**, 450
Goto, T., see Fukase, T., **263**, 420
Goto, T., see Seki, K., **263**, 290
Greven, M., see Endoh, Y., **263**, 349
Gu, G.D., T. Egi, N. Koshizuka, P.A. Miles, G.J. Russell and S.J. Kennedy, Effect of growth conditions on crystal morphology and superconductivity of Bi-2212 oxide, **263**, 180
Gu, G.D., see Han, S.H., **263**, 461
Gu, G.D., see Zhao, Y., **263**, 431
Gupta, A., see Tsuei, C.C., **263**, 232

Hagiwara, M., see Yamao, T., **263**, 464
Hahakura, S., see Ueyama, M., **263**, 172
Han, S.H., Y. Zhao, G.D. Gu, G.J. Russell and N. Koshizuka, Vortex-glass superconductivity of CuO_2 trilayers in BSCCO single crystals, **263**, 461
Han, S.H., see Maple, M.B., **263**, 490
Han, S.H., see Zhao, Y., **263**, 431
Hanaguri, T., T. Tsuboi, A. Maeda, T. Nishizaki, N. Kobayashi, Y. Kotaka, J.-i. Shimoyama and K. Kishio, Doping level dependence of magnetization anomalies and heat capacity of $Bi_2Sr_2CaCu_2O_{8+\delta}$ in the mixed state, **263**, 434
Hanaguri, T., see Maeda, A., **263**, 438
Harashina, H., K. Kodama, S. Shamoto, M. Sato, K. Kakurai and M. Nishi, Superconductivity-induced B_{2u} phonon anomalies of $YBa_2Cu_3O_{6+x}$ and symmetry of the order parameter. Neutron inelastic scattering studies, **263**, 257
Harashina, H., see Kodama, K., **263**, 333
Harris, J.M., see Tsui, O.K.C., **263**, 381
Hasegawa, M., see Maeda, A., **263**, 438
Hasegawa, T., see Kitazawa, K., **263**, 214
Hayashi, A., see Rykov, A.I., **263**, 359
Hayashi, K., see Ueyama, M., **263**, 172
Hayashi, N., see Machida, K., **263**, 428
Herrmann, J., see Maple, M.B., **263**, 490
Hervieu, M., see Raveau, B., **263**, 151
Hewat, A.W., see Khasanova, N.R., **263**, 200
Higuchi, T., S.I. Yoo, K. Sawada, N. Sakai and M. Murakami, Critical currents of OCMG processed Nd123 crystals, **263**, 396
Hikata, T., see Ueyama, M., **263**, 172
Hiraga, K., see Muraoka, Y., **263**, 193
Hirashima, S., see Itoh, M., **263**, 486
Honma, H., see Kijima, N., **263**, 523
Hosoya, S., see Endoh, Y., **263**, 349
Howard, C.J., see Takata, M., **263**, 176
Hussey, N.E., see Mackenzie, A.P., **263**, 510

Ichikawa, F., see Yamasaki, Y., **263**, 317
Ichioka, M., see Machida, K., **263**, 428
Ido, M., M. Oda, N. Momono, C. Manabe and T. Nakano, STM/STS and electronic specific heat of high-T_c cuprates. Symmetry of the order parameter, **263**, 225
Ido, M., see Oda, M., **263**, 241
Ihara, H., see Magishi, K., **263**, 375
Iino, Y. and M. Imada, Spin correlations in 1D Hubbard model at low temperatures and small dopings, **263**, 111
Iino, Y., see Maeda, A., **263**, 438
Ikebe, M., H. Fujishiro, K. Nakasato and K. Noto, Thermal conductivity of $REBa_2Cu_3O_7$ (RE = Y,Dy,Sm,Nd) superconducting oxides, **263**, 309
Ikebe, M., see Fujishiro, H., **263**, 305
Ikeda, S., K. Kumagai, J. Jiang, M.S. Lin, C.C. Lai and H.C. Ku, NMR study of quaternary superconductors RT_2B_2C (R = Y,La,Th, and T = Ni,Pd,Pt), **263**, 526
Ikeda, S., Y. Maeno, M. Nohara and T. Fujita, $Sr_2CaRu_2O_y$: a new phase of layered perovskite, **263**, 558
Ikeda, S., see Kato, M., **263**, 482
Imada, M., Fermi liquid phenomenology combined with scaling theory of the metal–insulator transition, **263**, 49
Imada, M., see Assaad, F.F., **263**, 78
Imada, M., see Iino, Y., **263**, 111
Imada, M., see Katoh, N., **263**, 114
Imada, M., see Yonemitsu, K., **263**, 82
Imazawa, C., see Takeuchi, S., **263**, 298
Inagaki, K., K. Yamaya and S. Tanda, Transport properties of localized states in $Bi_2Sr_2CuO_6$ single crystals, **263**, 321
Inoue, J. and S. Maekawa, Effects of Hund coupling on magneto-transport in perovskite Mn oxides, **263**, 138

Inoue, J., see Akazawa, S., **263**, 142
Inoue, J., see Ishihara, S., **263**, 130
Inoue, K., see Ishida, T., **263**, 260
Inoue, K., see Niimi, H., **263**, 412
Inoue, K., see Yamanaka, A., **263**, 516
Irie, A., see Oya, G.-i., **263**, 453
Ishida, K., Y. Kitaoka, K. Asayama, K. Kadowaki and T. Mochiku, Spin correlations in the normal state of a $Bi_2Sr_2CaCu_2O_8$ single crystal, **263**, 371
Ishida, T., K. Inoue, K. Okuda, H. Asaoka, Y. Kazumata, K. Noda and H. Takei, Anisotropy of superconductivity in an untwinned $YBa_2Cu_3O_7$ single crystal, **263**, 260
Ishida, T., see Nojima, T., **263**, 424
Ishihara, S., J. Inoue and S. Maekawa, Electronic structure and effective Hamiltonian in perovskite Mn oxides, **263**, 130
Ishikawa, Y., see Enomoto, Y., **263**, 21
Ishizuka, K., see Fukase, T., **263**, 420
Itoh, M., M. Sugahara, S. Hirashima, T. Yamauchi and Y. Ueda, Microscopic magnetic property and electronic state of the spin-Peierls compound $CuGeO_3$: $^{63,65}Cu$ and ^{17}O NMR studies, **263**, 486
Itoh, Y., Quantum ground states and spin-fluctuation spectrum in high-T_c Cu oxides, **263**, 378
Itti, R., see Ting, W., **263**, 185
Iyo, A., see Magishi, K., **263**, 375
Izumi, F., see Khasanova, N.R., **263**, 200

Jiang, J., see Ikeda, S., **263**, 526
Julian, S.R., see Mackenzie, A.P., **263**, 510

Kadowaki, K., Sudden disappearance of c-axis dissipation in single-crystalline $Bi_2Sr_2CaCu_2O_{8+\delta}$, **263**, 164
Kadowaki, K., see Ishida, K., **263**, 371
Kadowaki, K., see Matsuda, Y., **263**, 457
Kaga, H., see Saikawa, T., **263**, 74
Kajimura, K., see Tanaka, Y., **263**, 238
Kajitani, T., see Takeuchi, S., **263**, 298
Kakurai, K., see Harashina, H., **263**, 257
Kakurai, K., see Kodama, K., **263**, 333
Kamata, M., see Fukase, T., **263**, 420
Kamimura, H., S. Matsuno, Y. Suwa and H. Ushio, A new theory of high-temperature superconductivity in cuprates, **263**, 126
Kaper, H.G., see Crabtree, G.W., **263**, 401
Karppinen, M., see Yamauchi, H., **263**, 146
Kasai, M., see Tokura, Y., **263**, 544
Kasai, M., see Yokoya, T., **263**, 505
Kashiwaya, S., see Tanaka, Y., **263**, 238
Kastner, M.A., see Endoh, Y., **263**, 349
Katano, S., see Nakata, H., **263**, 344
Katano, S., see Takeuchi, S., **263**, 298
Katayama-Yoshida, H., see Yokoya, T., **263**, 505
Kato, M., K. Shiota, S. Ikeda, Y. Maeno, T. Fujita and Y. Koike, Transport and magnetic properties of the spin-$\frac{1}{2}$ ladder system $(Sr,A)_{14}Cu_{24}O_{41}$ (A = Ba and Ca), **263**, 482
Kato, M., see Takeuchi, S., **263**, 298
Katoh, N. and M. Imada, Spin gap in generalized spin models with dimerization and the model for CaV_4O_9, **263**, 114
Kawachi, M., see Yamao, T., **263**, 464
Kawamata, S., see Niimi, H., **263**, 412
Kawamura, H., see Fujiyoshi, T., **263**, 446
Kazumata, Y., see Ishida, T., **263**, 260
Kennedy, S.J., see Gu, G.D., **263**, 180
Ketchen, M.B., see Tsuei, C.C., **263**, 232
Khasanova, N.R., F. Izumi, E. Takayama-Muromachi and A.W. Hewat, Crystal structure of the superconductor $Ca_2Sr_2Cu_3GaO_9$ prepared at high pressure, **263**, 200
Kichambare, P.D., see Kijima, N., **263**, 523
Kijima, N., P.D. Kichambare, H. Honma, S. Ebisu and S. Nagata, Superconductivity in $Li(Ti_{1-x}V_x)_2O_4$, **263**, 523
Kikuchi, M., M. Shida, E. Ohshima, Y. Muraoka and Y. Syono, Synthesis conditions and superconductivity of $Pb_{0.7}Cu_{0.3}Sr_2Y_{0.5}Ca_{0.5}Cu_2O_y$, **263**, 204
Kikuchi, M., see Goto, T., **263**, 253
Kikuchi, M., see Muraoka, Y., **263**, 193
Kikuchi, M., see Ohshima, E., **263**, 189
Kirtley, J.R., see Tsuei, C.C., **263**, 232
Kishine, J.-I., Spin fluctuations in an itinerant Heisenberg system. Naive RPA and vertex correction, **263**, 99
Kishio, K., see Hanaguri, T., **263**, 434
Kitamori, N., see Sugai, S., **263**, 363
Kitamura, Y., see Koyama, K., **263**, 336
Kitaoka, Y., see Akimitsu, J., **263**, 475
Kitaoka, Y., see Ishida, K., **263**, 371
Kitaoka, Y., see Magishi, K., **263**, 375
Kitaoka, Y., see Zheng, G.-q., **263**, 367
Kitazawa, K., H. Sugawara and T. Hasegawa, On the superconducting gap structure of high-temperature superconductors by STM/STS, **263**, 214
Kivelson, S.A., see Emery, V.J., **263**, 44
Kluge, T., see Koike, Y., **263**, 329
Kobayashi, N., see Hanaguri, T., **263**, 434
Kobayashi, N., see Koike, Y., **263**, 329
Kobayashi, N., see Muraoka, Y., **263**, 193
Kobayashi, N., see Noji, T., **263**, 442
Kobayashi, N., see Onodera, Y., **263**, 409
Kobayashi, S., see Ueyama, M., **263**, 172
Kobayashi, Y., see Nishikawa, T., **263**, 554
Kodama, K., S. Shamoto, H. Harashina, M. Sato, M. Nishi and K. Kakurai, T_c-suppression mechanism of Pr-doping in (R, Pr)$Ba_2Cu_3O_7$, **263**, 333
Kodama, K., see Harashina, H., **263**, 257
Kodama, K., see Takata, M., **263**, 176
Kodama, K., see Takata, M., **263**, 340
Kodama, M., see Ekino, T., **263**, 249
Kodama, M., see Nakamura, F., **263**, 282
Kodama, Y., see Zheng, G.-q., **263**, 367
Koike, Y., A. Fujiwara, T. Kluge, T. Noji, T. Nishizaki, N. Kobayashi, A. Yamanaka, F. Minami and S. Takekawa, Dimensionality, T_c and Cu-site substitution effect of iodine-intercalated and oxidized $Bi_2Sr_2CaCu_2O_{8+\delta}$: Interpretation by the multilayer model, **263**, 329
Koike, Y., see Kato, M., **263**, 482

Koike, Y., see Noji, T., **263**, 442
Koike, Y., see Sugai, S., **263**, 363
Koike, Y., see Takeuchi, S., **263**, 298
Kondo, R., see Yasuda, T., **263**, 416
Koshelev, A.E., see Crabtree, G.W., **263**, 401
Koshibae, W., Y. Ohta and S. Maekawa, Dynamics of a hole in spin 1 chain, **263**, 118
Koshizuka, N., see Gu, G.D., **263**, 180
Koshizuka, N., see Han, S.H., **263**, 461
Koshizuka, N., see Ting, W., **263**, 185
Koshizuka, N., see Zhao, Y., **263**, 431
Kotaka, Y., see Hanaguri, T., **263**, 434
Koyama, K., T. Tange, Y. Kitamura, T. Saito and K. Mizuno, Oxygen concentration dependence of superconductivity in Pr-substituted $Ln_{1-x}Pr_xBa_2Cu_3O_{7-y}$ (Ln = Y, Nd, Gd), **263**, 336
Koyama, T. and M. Tachiki, The effect of internal orbital motion in 2D d-wave superconductors, **263**, 25
Koyama, T., see Tachiki, M., **263**, 1
Koyanagi, M., see Tanaka, Y., **263**, 238
Krishana, K., see Tsui, O.K.C., **263**, 381
Ku, H.C., see Ikeda, S., **263**, 526
Kubo, S., see Yamasaki, Y., **263**, 317
Kubota, H., see Fujiyoshi, T., **263**, 446
Kumagai, K., see Ikeda, S., **263**, 526
Kumagai, K., see Matsuda, Y., **263**, 457
Kuroda, K., see Ting, W., **263**, 185
Kuwahara, H., see Tokura, Y., **263**, 544
Kuwasawa, Y., see Nojima, T., **263**, 424
Kwok, W.K., see Crabtree, G.W., **263**, 401

Lai, C.C., see Ikeda, S., **263**, 526
Lang, M., see Steglich, F., **263**, 498
Lang, M., see Takahashi, S., **263**, 30
Lanza, H., see Civale, L., **263**, 389
Leaf, G.K., see Crabtree, G.W., **263**, 401
Lee, C.H., see Endoh, Y., **263**, 349
Levine, D.M., see Crabtree, G.W., **263**, 401
Levy, P., see Civale, L., **263**, 389
Liang, W.Y., J.W. Loram, K.A. Mirza, N. Athanassopoulou and J.R. Cooper, Specific heat and susceptibility determination of the pseudogap in $YBCO_{7-\delta}$, **263**, 277
Lin, M.S., see Ikeda, S., **263**, 526
Locquet, J.-P., Local electrochemical oxidation/reduction: first step towards a new lithography?, **263**, 160
Loeser, A.G., D.S. Dessau and Z.-X. Shen, Doping dependence of $Bi_2Sr_2CaCu_2O_{8+\delta}$ in the normal state, **263**, 208
Lonzarich, G.G., see Mackenzie, A.P., **263**, 510
Loram, J.W., see Liang, W.Y., **263**, 277
Lühmann, T., see Steglich, F., **263**, 498

Machida, K., M. Ichioka, N. Hayashi and N. Enomoto, Vortex core structure and possible pairing mixture in d-wave superconductors, **263**, 428
Mackenzie, A.P., S.R. Julian, A.J. Diver, G.G. Lonzarich, N.E. Hussey, Y. Maeno, S. Nishizaki and T. Fujita, Calculation of thermodynamic and transport properties of Sr_2RuO_4 at low temperatures using known Fermi surface parameters, **263**, 510
Maeda, A., T. Hanaguri, Y. Iino, S. Masuoka, Y. Matsushita, M. Hasegawa and H. Takei, Nonlinear Meissner effect of the cuprate superconductors investigated by London penetration depth measurement, **263**, 438
Maeda, A., see Hanaguri, T., **263**, 434
Maekawa, S., T. Tohyama and S. Yunoki, Anomalous low-energy excitations in strongly correlated electron systems, **263**, 61
Maekawa, S., see Akazawa, S., **263**, 142
Maekawa, S., see Enomoto, Y., **263**, 21
Maekawa, S., see Inoue, J., **263**, 138
Maekawa, S., see Ishihara, S., **263**, 130
Maekawa, S., see Koshibae, W., **263**, 118
Maekawa, S., see Ohta, Y., **263**, 94
Maekawa, S., see Tsutsui, K., **263**, 107
Maekawa, S., see Yunoki, S., **263**, 90
Maeno, Y., see Ikeda, S., **263**, 558
Maeno, Y., see Kato, M., **263**, 482
Maeno, Y., see Mackenzie, A.P., **263**, 510
Maeno, Y., see Nakamura, F., **263**, 282
Maeno, Y., see Yamanaka, A., **263**, 516
Maeno, Y., see Yoshida, K., **263**, 519
Magishi, K., G.-q. Zheng, Y. Kitaoka, K. Asayama, K. Tokiwa, A. Iyo and H. Ihara, ^{63}Cu NMR investigation of $HgBa_2Ca_2Cu_3O_{8+\delta}$, **263**, 375
Manabe, C., see Ido, M., **263**, 225
Manabe, C., see Oda, M., **263**, 241
Mancini, F., S. Marra and H. Matsumoto, Magnetic properties of the two-dimensional Hubbard model, **263**, 70
Mancini, F., see Matsumoto, H., **263**, 66
Maple, M.B., M.C. de Andrade, J. Herrmann, S.H. Han, R. Movshovich, D.A. Gajewski and R. Chau, Non-Fermi liquid behavior in chemically-substituted f-electron compounds, **263**, 490
Marra, S., see Mancini, F., **263**, 70
Marra, S., see Matsumoto, H., **263**, 66
Masuoka, S., see Maeda, A., **263**, 438
Matsuda, A., see Watanabe, T., **263**, 313
Matsuda, Y., M.B. Gaifullin, K. Kumagai, K. Kadowaki and T. Mochiku, Microwave response in the vortex state of $Bi_2Sr_2CaCu_2O_{8+\delta}$, **263**, 457
Matsumoto, H., S. Odashima, F. Mancini and S. Marra, Effects of two-site local correlation in the 2D Hubbard model, **263**, 66
Matsumoto, H., see Mancini, F., **263**, 70
Matsumoto, S., see Akimitsu, J., **263**, 475
Matsuno, S., see Kamimura, H., **263**, 126
Matsushita, Y., see Maeda, A., **263**, 438
Matsuura, M., see Yamao, T., **263**, 464
Matsuyama, A., T. Sasaki, T. Fukase and N. Toyota, Hall effect in the mixed state of the organic superconductor κ-(BEDT-TTF)$_2$Cu(NCS)$_2$, **263**, 534
Matsuyama, A., see Sasaki, T., **263**, 538
Michel, C., see Raveau, B., **263**, 151
Miles, P.A., see Gu, G.D., **263**, 180

Minami, F., see Koike, Y., **263**, 329
Mirza, K.A., see Liang, W.Y., **263**, 277
Miyahara, K., see Fujiyoshi, T., **263**, 446
Mizuhashi, K., see Zheng, G.-q., **263**, 367
Mizukoshi, T., see Yao, X., **263**, 197
Mizuno, K., see Koyama, K., **263**, 336
Mizuno, Y., see Yunoki, S., **263**, 90
Mochiku, T., see Ishida, K., **263**, 371
Mochiku, T., see Matsuda, Y., **263**, 457
Modler, R., see Steglich, F., **263**, 498
Modler, R., see Takahashi, S., **263**, 30
Momono, N., see Ido, M., **263**, 225
Môri, N., see Akimitsu, J., **263**, 475
Morioka, Y., see Muraoka, Y., **263**, 193
Moritomo, Y., see Tokura, Y., **263**, 544
Movshovich, R., see Maple, M.B., **263**, 490
Murakami, M., see Higuchi, T., **263**, 396
Muranaka, K., see Ueyama, M., **263**, 172
Muraoka, Y., N. Ohnishi, M. Nagoshi, Y. Morioka, M. Kikuchi, K. Hiraga, N. Kobayashi and Y. Syono, AgI-intercalation into $Bi_2Sr_2CaCu_2O_z$ compound, **263**, 193
Muraoka, Y., see Kikuchi, M., **263**, 204
Murase, M., see Ohta, Y., **263**, 94
Murayama, Y., see Sugano, R., **263**, 17

Nagaosa, N., Interlayer charge dynamics of high-T_c cuprates, **263**, 54
Nagata, S., see Kijima, N., **263**, 523
Nagata, T., see Akimitsu, J., **263**, 475
Nagoshi, M., Y. Syono, M. Tachiki and Y. Fukuda, Core-level binding energy as a probe for charge distributions in high-T_c cuprates, **263**, 294
Nagoshi, M., see Muraoka, Y., **263**, 193
Nagoshi, M., see Ohshima, E., **263**, 189
Nagoshi, M., see Sanada, N., **263**, 286
Nakadaira, T., see Sanada, N., **263**, 286
Nakajima, S., see Goto, T., **263**, 253
Nakamura, F., M. Kodama, S. Sakita, Y. Maeno and T. Fujita, Transport properties of detwinned crystals of $La_{2-x}Sr_xCuO_4$, **263**, 282
Nakamura, F., see Ekino, T., **263**, 249
Nakano, T., see Ido, M., **263**, 225
Nakasato, K., see Fujishiro, H., **263**, 305
Nakasato, K., see Ikebe, M., **263**, 309
Nakata, H., H. Okajima, T. Yokoo, A. Yamashita, J. Akimitsu and S. Katano, The phase diagram of the new superconductor $Sr_2CuO_2(CO_3)_{1-x}(BO_3)_x$, **263**, 344
Nelson, D.R., Points, lines and planes: vortex pinning in high-temperature superconductors, **263**, 12
Nieva, G., see Civale, L., **263**, 389
Niimi, H., K. Inoue, S. Kawamata and K. Okuda, Study of magnetic flux in a Bi2212 single crystal by torsional oscillation, **263**, 412
Nishi, M., see Harashina, H., **263**, 257
Nishi, M., see Kodama, K., **263**, 333
Nishibori, E., see Takata, M., **263**, 176
Nishikawa, T., Y. Yasui, Y. Kobayashi and M. Sato, Studies of the metal–insulator transition of misfit-layer compounds $La_{1.17-x}Sr_xVS_{3.17}$, **263**, 554
Nishizaki, S., see Mackenzie, A.P., **263**, 510
Nishizaki, S., see Yamanaka, A., **263**, 516
Nishizaki, S., see Yoshida, K., **263**, 519
Nishizaki, T., see Hanaguri, T., **263**, 434
Nishizaki, T., see Koike, Y., **263**, 329
Nishizaki, T., see Noji, T., **263**, 442
Nishizaki, T., see Onodera, Y., **263**, 409
Nishizaki, T., see Yamasaki, Y., **263**, 317
Noda, K., see Ishida, T., **263**, 260
Nohara, M., see Ikeda, S., **263**, 558
Noji, T., T. Takabayashi, T. Nishizaki, N. Kobayashi and Y. Koike, Irreversibility line and second peak in the magnetization curves of $Pb_2Sr_2Y_{1-x}Ca_xCu_3O_8$ single crystals, **263**, 442
Noji, T., see Koike, Y., **263**, 329
Noji, T., see Sugai, S., **263**, 363
Nojima, T., T. Ishida and Y. Kuwasawa, Current–voltage characteristics of $YBa_2Cu_3O_{7-\delta}$ films at low magnetic fields, **263**, 424
Noto, K., see Fujishiro, H., **263**, 305
Noto, K., see Ikebe, M., **263**, 309

Oda, M., C. Manabe, T. Abe and M. Ido, STM and STS studies on the superconducting gap symmetry of $Bi_2Sr_2CaCu_2O_{8+\delta}$, **263**, 241
Oda, M., see Ido, M., **263**, 225
Odaguchi, T., see Zheng, G.-q., **263**, 367
Odashima, S., see Matsumoto, H., **263**, 66
Ohkura, K., see Ueyama, M., **263**, 172
Ohmori, C., see Saikawa, T., **263**, 74
Ohnishi, N., see Muraoka, Y., **263**, 193
Ohshima, E., M. Kikuchi, M. Nagoshi and Y. Syono, Cation substitution and deoxygenation effect on superconductivity of Tl–Sr1212 cuprate, **263**, 189
Ohshima, E., see Kikuchi, M., **263**, 204
Ohta, Y., M. Murase, R. Eder and S. Maekawa, Impurity effect in the 2D t–J model, **263**, 94
Ohta, Y., see Eder, R., **263**, 104
Ohta, Y., see Koshibae, W., **263**, 118
Ohta, Y., see Tsutsui, K., **263**, 107
Okajima, H., see Nakata, H., **263**, 344
Okuda, K., see Ishida, T., **263**, 260
Okuda, K., see Niimi, H., **263**, 412
Ong, N.P., see Tsui, O.K.C., **263**, 381
Ono, Y., see Takeuchi, S., **263**, 298
Onodera, Y., T. Nishizaki, N. Kobayashi, H. Asaoka and H. Takei, Magnetization jump due to the first-order vortex lattice melting transition in $YBa_2Cu_3O_y$ single crystals, **263**, 409
Onogi, T., see Sugano, R., **263**, 17
Ōnuki, Y., see Steglich, F., **263**, 498
Oya, G.-i. and A. Irie, Fluxon motion in the stacked intrinsic Josephson junctions in $Bi_2Sr_2CaCu_2O_y$ single crystals, **263**, 453

Pasquini, G., see Civale, L., **263**, 389
Paulsen, C., see Steglich, F., **263**, 498
Peterson, J.B., see Tsui, O.K.C., **263**, 381

Racah, D. and G. Deutscher, Properties of normal metal/dielectric/high-T_c junctions obtained by in-situ oxidation, **263**, 218
Raveau, B., M. Hervieu and C. Michel, Generation of superconducting oxycarbonates and layered cuprates through shearing mechanisms, **263**, 151
Riera, J., see Tsutsui, K., **263**, 107
Rupp, M., see Tsuei, C.C., **263**, 232
Russell, G.J., see Gu, G.D., **263**, 180
Russell, G.J., see Han, S.H., **263**, 461
Russell, G.J., see Zhao, Y., **263**, 431
Rykov, A.I., Y. Ueda and A. Hayashi, Magnetic susceptibility of the orthorhombic 2D-antiferromagnet $YSr_2Cu_2GaO_7$, **263**, 359

Saga, N., see Ueyama, M., **263**, 172
Saikawa, T., C. Ohmori, H. Kaga and P. Fulde, Renormalizations of the Landau Fermi-liquid parameter in the Hubbard model and the three-band model, **263**, 74
Saito, T., see Koyama, K., **263**, 336
Sakai, N., see Higuchi, T., **263**, 396
Sakata, M., see Takata, M., **263**, 176
Sakata, M., see Takata, M., **263**, 340
Sakita, S., see Ekino, T., **263**, 249
Sakita, S., see Nakamura, F., **263**, 282
Salk, S.-H.S., see Doh, H., **263**, 86
Sanada, N., T. Nakadaira, M. Shimomura, Y. Suzuki, Y. Fukuda, M. Nagoshi, Y. Syono and M. Tachiki, Electronic states and superconducting properties of $Bi_2Sr_2(Ca, Gd)Cu_2O_y$ studied by X-ray photoelectron and inverse-photoemission spectroscopy, **263**, 286
Sasada, M., see Yamanaka, A., **263**, 516
Sasaki, S., see Takata, M., **263**, 340
Sasaki, T., A. Matsuyama and T. Fukase, Resistive jump and hysteresis in the vortex state of the organic superconductor κ-(BEDT-TTF)$_2$Cu(NCS)$_2$, **263**, 538
Sasaki, T., see Fukase, T., **263**, 420
Sasaki, T., see Matsuyama, A., **263**, 534
Sato, H., see Fukushima, K., **263**, 122
Sato, K., see Ueyama, M., **263**, 172
Sato, M., Anomalous metallic phase in Cu oxides and other 2D Mott systems, **263**, 271
Sato, M., see Harashina, H., **263**, 257
Sato, M., see Kodama, K., **263**, 333
Sato, M., see Nishikawa, T., **263**, 554
Sato, M., see Shamoto, S., **263**, 550
Sato, M., see Takata, M., **263**, 176
Sato, M., see Takata, M., **263**, 340
Sato, M., see Tanaka, S., **263**, 245
Sato, Y., see Terasaki, I., **263**, 302
Sawada, K., see Higuchi, T., **263**, 396
Seki, K., T. Goto and T. Fukase, Ultrasonic and La-NMR study on structural phase transitions and antiferromagnetic ordering in $La_{2-x-y}Ba_xCe_yCu_{1-z}Zn_zO_4$ ($x \cong 1/8$), **263**, 290
Shamoto, S., S. Tanaka, E. Ueda and M. Sato, Single crystal growth of $BaCo_{1-x}Ni_xS_2$, **263**, 550
Shamoto, S., see Harashina, H., **263**, 257
Shamoto, S., see Kodama, K., **263**, 333
Shen, Z.-X., see Loeser, A.G., **263**, 208
Shibauchi, T., see Tamegai, T., **263**, 325
Shida, M., see Kikuchi, M., **263**, 204
Shimomura, M., see Sanada, N., **263**, 286
Shimoyama, J.-i., see Hanaguri, T., **263**, 434
Shimura, T., see Tamegai, T., **263**, 325
Shiohara, Y., see Yao, X., **263**, 197
Shiota, K., see Kato, M., **263**, 482
Shirane, G., see Endoh, Y., **263**, 349
Steglich, F., R. Modler, P. Gegenwart, M. Deppe, M. Weiden, M. Lang, C. Geibel, T. Lühmann, C. Paulsen, J.L. Tholence, Y. Ōnuki, M. Tachiki and S. Takahashi, Experimental evidence for a generalized FFLO state in clean type-II superconductors with short coherence length and enhanced Pauli susceptibility, **263**, 498
Steglich, F., see Takahashi, S., **263**, 30
Sueyoshi, T., see Fujiyoshi, T., **263**, 446
Sugahara, M., see Itoh, M., **263**, 486
Sugai, S., N. Kitamori, N. Yamane, T. Noji and Y. Koike, Phonon and two-magnon Raman scattering in $Pb_2Sr_2Ho(Y)_{1-x}Ca_xCu_3O_{8+\delta}$, **263**, 363
Sugano, R., T. Onogi and Y. Murayama, Monte Carlo study of vortex-line dynamics with correlated disorders, **263**, 17
Sugawara, H., see Kitazawa, K., **263**, 214
Sun, J.Z., see Tsuei, C.C., **263**, 232
Suwa, Y., see Kamimura, H., **263**, 126
Suzuki, J., see Tamegai, T., **263**, 325
Suzuki, M., see Yamasaki, Y., **263**, 317
Suzuki, S., see Yamamoto, T., **263**, 530
Suzuki, T., see Fukase, T., **263**, 420
Suzuki, Y., see Sanada, N., **263**, 286
Syono, Y., see Goto, T., **263**, 253
Syono, Y., see Kikuchi, M., **263**, 204
Syono, Y., see Muraoka, Y., **263**, 193
Syono, Y., see Nagoshi, M., **263**, 294
Syono, Y., see Ohshima, E., **263**, 189
Syono, Y., see Sanada, N., **263**, 286

Tachiki, M., T. Koyama and S. Takahashi, Some exotic phenomena related to vortices in high-T_c and heavy fermion superconductors, **263**, 1
Tachiki, M., see Koyama, T., **263**, 25
Tachiki, M., see Nagoshi, M., **263**, 294
Tachiki, M., see Sanada, N., **263**, 286
Tachiki, M., see Steglich, F., **263**, 498
Tachiki, M., see Takahashi, S., **263**, 30
Tajima, S., see Terasaki, I., **263**, 302
Takabayashi, T., see Noji, T., **263**, 442
Takahashi, H., see Akimitsu, J., **263**, 475
Takahashi, K., see Yamamoto, T., **263**, 530
Takahashi, S., M. Tachiki, R. Modler, P. Gegenwart, M. Lang and F. Steglich, Theory of a generalized Fulde–Ferrell–Larkin–Ovchinnikov state in heavy fermion and intermediate-valence superconductors, **263**, 30

Takahashi, S., see Steglich, F., **263**, 498
Takahashi, S., see Tachiki, M., **263**, 1
Takahashi, T., see Yokoya, T., **263**, 505
Takano, M., Spin ladder compounds, **263**, 468
Takano, S., see Yasuda, T., **263**, 416
Takata, M., E. Nishibori, T. Takayama, M. Sakata, K. Kodama, M. Sato and C.J. Howard, The nuclear density of YBCO by the maximum entropy method using neutron powder data, **263**, 176
Takata, M., T. Takayama, M. Sakata, S. Sasaki, K. Kodama and M. Sato, Direct observation of Pr–CuO_2 hybridization in $PrBa_2Cu_3O_{7-\delta}$ by the MEM charge density study, **263**, 340
Takayama-Muromachi, E., see Khasanova, N.R., **263**, 200
Takayama, T., see Takata, M., **263**, 176
Takayama, T., see Takata, M., **263**, 340
Takei, H., see Ishida, T., **263**, 260
Takei, H., see Maeda, A., **263**, 438
Takei, H., see Onodera, Y., **263**, 409
Takekawa, S., see Koike, Y., **263**, 329
Takenaka, K., see Uchida, S., **263**, 264
Takeuchi, S., C. Imazawa, S. Katano, M. Kato, Y. Ono, T. Kajitani and Y. Koike, Correlation between superconductivity, electronic state and crystal structure in $La_{2-y-x}Bi_yBa_xCuO_4$, **263**, 298
Tamasaku, K., see Tanaka, S., **263**, 245
Tamasaku, K., see Uchida, S., **263**, 264
Tamegai, T., T. Shimura, J. Suzuki and T. Shibauchi, Transport properties of metallic cuprates $La_6BaLnCu_8O_{20}$ (Ln: lanthanide), **263**, 325
Tanaka, S., E. Ueda, M. Sato, K. Tamasaku and S.-i. Uchida, Anisotropy of the energy gap structure of high-T_c oxides observed by STM/STS, **263**, 245
Tanaka, S., see Shamoto, S., **263**, 550
Tanaka, S., see Terasaki, I., **263**, 302
Tanaka, S., see Ting, W., **263**, 185
Tanaka, S., see Yamauchi, H., **263**, 146
Tanaka, Y., S. Kashiwaya, M. Koyanagi and K. Kajimura, Tunneling effects in d-wave superconductors, **263**, 238
Tanda, S., see Inagaki, K., **263**, 321
Tange, T., see Koyama, K., **263**, 336
Terasaki, I., Y. Sato, S. Tajima and S. Tanaka, Anisotropic magnetoresistance in slightly overdoped $YBa_2Cu_3O_y$ crystals, **263**, 302
Tholence, J.L., see Steglich, F., **263**, 498
Ting, W., T. Egi, R. Itti, K. Kuroda, N. Koshizuka and S. Tanaka, STM/STS measurements on clean and stable surfaces of as-prepared $Nd_1Ba_2Cu_3O_y$ single crystals, **263**, 185
Tohyama, T., see Maekawa, S., **263**, 61
Tokiwa, K., see Magishi, K., **263**, 375
Tokura, Y., Y. Tomioka, H. Kuwahara, A. Asamitsu, Y. Moritomo and M. Kasai, Magnetic-field-induced insulator–metal phenomena in perovskite manganese oxides, **263**, 544
Tokura, Y., see Yokoya, T., **263**, 505
Tomioka, Y., see Tokura, Y., **263**, 544
Toyota, N., see Matsuyama, A., **263**, 534
Tsuboi, T., see Hanaguri, T., **263**, 434
Tsuei, C.C., J.R. Kirtley, M. Rupp, J.Z. Sun, C.C. Chi, A. Gupta, L.S. Yu-Jahnes and M.B. Ketchen, Half-integer flux quantum effect in tricrystal cuprate superconductors, **263**, 232
Tsui, O.K.C., K. Krishana, J.M. Harris, N.P. Ong and J.B. Peterson, The Josephson plasma mode in $Bi_2Sr_2CaCu_2O_8$ and quasi-particle mean-free-path in $YBa_2Cu_3O_7$, **263**, 381
Tsutsui, K., Y. Ohta, R. Eder, S. Maekawa, E. Dagotto and J. Riera, Low energy excitations of the Anderson lattice model, **263**, 107

Uchida, S., Y. Fukuzumi, K. Takenaka and K. Tamasaku, Zn-doping effects on the charge dynamics in high-T_c copper oxides. Spin-gap effect and Josephson plasma in the underdoped regime, **263**, 264
Uchida, S., see Zheng, G.-q., **263**, 367
Uchida, S.-i., see Tanaka, S., **263**, 245
Udagawa, M., see Yamanaka, A., **263**, 516
Ueda, E., see Shamoto, S., **263**, 550
Ueda, E., see Tanaka, S., **263**, 245
Ueda, Y., see Itoh, M., **263**, 486
Ueda, Y., see Rykov, A.I., **263**, 359
Uehara, M., see Akimitsu, J., **263**, 475
Ueyama, M., K. Ohkura, K. Hayashi, S. Kobayashi, K. Muranaka, T. Hikata, N. Saga, S. Hahakura and K. Sato, Development of Ag-sheathed Bi(2223) superconducting wires and coils, **263**, 172
Ushio, H., see Kamimura, H., **263**, 126

Varma, C.M., Theory of the Cu–O metals, **263**, 39
Vinokur, V.M., see Crabtree, G.W., **263**, 401

Wakimoto, S., see Endoh, Y., **263**, 349
Watanabe, T. and A. Matsuda, Magnetoresistance and high-temperature resistivity of $Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_2O_{8+\delta}$ single crystals, **263**, 313
Weiden, M., see Steglich, F., **263**, 498

Yamada, K., see Endoh, Y., **263**, 349
Yamamoto, T., S. Suzuki, K. Takahashi and Y. Yoshisato, Analysis of the tunneling characteristics of $Ba_{1-x}K_xBiO_3$ superconducting thin films based on the gap-energy distribution model, **263**, 530
Yamanaka, A., N. Asayama, M. Sasada, K. Inoue, M. Udagawa, S. Nishizaki, Y. Maeno and T. Fujita, Electronic Raman scattering spectra of Sr_2RuO_4 single crystals, **263**, 516
Yamanaka, A., see Koike, Y., **263**, 329
Yamane, N., see Sugai, S., **263**, 363
Yamao, T., M. Kawachi, M. Hagiwara and M. Matsuura, Successive transitions of a hierarchical nature in a superconductive ceramic of $YBa_2Cu_4O_8$. Electrical resistivity, **263**, 464
Yamasaki, Y., T. Fukami, T. Nishizaki, F. Ichikawa, T. Aomine, S. Kubo and M. Suzuki, Magnetoresistance of superconducting $Nd_{2-x}Ce_xCuO_4$ in magnetic fields parallel to the c-axis, **263**, 317
Yamashita, A., see Nakata, H., **263**, 344
Yamauchi, H., M. Karppinen and S. Tanaka, Homologous series of layered cuprates, **263**, 146
Yamauchi, T., see Itoh, M., **263**, 486
Yamaya, K., see Inagaki, K., **263**, 321

Yao, X., T. Mizukoshi, M. Egami and Y. Shiohara, Superconductivity and crystal growth of YBCO grown under 1 atm oxygen pressure, **263**, 197

Yasuda, T., R. Kondo, S. Takano, T. Fukami and T. Aomine, Interlayer Josephson coupling in Bi–Sr–Ca–Cu–O single crystals, **263**, 416

Yasui, Y., see Nishikawa, T., **263**, 554

Yokoo, T., see Nakata, H., **263**, 344

Yokoya, T., A. Chainani, T. Takahashi, H. Katayama-Yoshida, M. Kasai and Y. Tokura, Extended van-Hove singularity in Sr_2RuO_4 observed by angle-resolved photoemission, **263**, 505

Yonemitsu, K. and M. Imada, Electron–phonon interaction and spin gap in nearly-half-filled correlated electron systems, **263**, 82

Yoo, S.I., see Higuchi, T., **263**, 396

Yoshida, K., Y. Maeno, S. Nishizaki and T. Fujita, Anisotropic superconductivity of Sr_2RuO_4, **263**, 519

Yoshisato, Y., see Yamamoto, T., **263**, 530

Yu-Jahnes, L.S., see Tsuei, C.C., **263**, 232

Yunoki, S., Y. Mizuno and S. Maekawa, Quantum Monte Carlo study of a magnetic impurity in a strongly correlated electron system, **263**, 90

Yunoki, S., see Maekawa, S., **263**, 61

Zhao, Y., G.D. Gu, S.H. Han, G.J. Russell and N. Koshizuka, Asymptotic power spectrum analysis of electrical noise in the vortex state of $Bi_2Sr_2CaCu_2O_{8+x}$ single crystals, **263**, 431

Zhao, Y., see Han, S.H., **263**, 461

Zheng, G.-q., T. Odaguchi, Y. Kitaoka, K. Asayama, Y. Kodama, K. Mizuhashi and S. Uchida, Zn and Ni doping effects on T_c and spin gap behavior in $YBa_2Cu_4O_8$ and $YBa_2Cu_3O_{6.6}$, **263**, 367

Zheng, G.-q., see Magishi, K., **263**, 375

ELSEVIER

Physica C 263 (1996) 571–573

PHYSICA C

Analytic Subject Index to Volume 263

AGI intercalation
–in $Bi_2Sr_2CaCu_2O_z$ 193
Anderson lattice model 107
Anisotropic order parameter 245
Anisotropic superconductor 438
Anisotropy
–of $Bi_2Sr_2CaCu_2O_8$ 381
–of Sr_2RuO_4 519
Anomalous metallic phase 271
Anomalous normal state 39
Anomaly 363
Antiferromagnetic order 86
–in $YSr_2Cu_2GaO_7$ 359
Atomic Thermal Vibration 176

C-axis resistivity
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystal 164
Charge density 340
Charge order 44
Charge transfer 294
–in $TlSr_2Ca_{1-x}Y_xCu_2O_y$ 189
Columnar defects 17
–in $YBa_2Cu_3O_{7-x}$ single crystals 389
Core level 294
Coulomb interactions 122
Critical current density
–in $Bi_2Sr_2CaCu_2O_{8+x}$ single crystals 416
–of Nd123 crystals 396
–of Y123 filaments 450
Critical current 12
Critical phenomena 49
Crystal growth
–of $Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_{2.0}O_x$ 180
–of YBCO 197
Current-voltage characteristics
–of $YBa_2Cu_3O_{7-\delta}$ film 424

D-wave superconductor 25, 225, 238, 428
D-wave 126
d-wave 253
Detwinning
–in $La_{2-x}Sr_xCuO_4$ 282
Dimensionality 329
Double-exchange model 134

Effective Hamiltonian
–in Mn oxides 130
Eilenberger theory 428
E-J characteristics 446
Electrical resistivity
–of $(Sr_{0.4}Ca_{0.6})_{14}Cu_{24-x}Co_xO_{41-\delta}$ 475
–of κ-(BEDT-TTF)$_2$Cu(NCS)$_2$ 538
–of $La_{2-x}Sr_xCuO_4$ 282
–of $La_{6-x}Sr_xBaLnCu_8O_{20}$ 325
–of $Li(Ti_{1-x}V_x)_2O_4$ 523
–of $Pb_{0.7}Cu_{0.3}Sr_2Y_{0.5}Cu_2O_y$ 204
–of $Sr_{14-x}Ba_xCu_{24}O_{41}$ 482
–of $Sr_{14-x}Ca_xCu_{24}O_{41}$ 482
–of $YBa_2Cu_3O_7$ 401
–of $YBa_2Cu_4O_8$ 464
Electron-doped superconductor 355
Electron-electron scattering 313
Electron-phonon interaction 82
Electronic specific heat
–of $La_{2-x}Sr_xCuO_4$ 225
Electronic state 66
Electronic structure 208
–in Mn oxides 130
–of Sr_2RuO_4 505
Energy gap
–of $La_{1.85}Sr_{0.15}CuO_4$ single crystals 249
Exact diagonalization 107
Excitations 61

FFLO state 1, 30, 498
Fermi liquid state 35
Fermi surface
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 208
–of Sr_2RuO_4 505
–of Sr_2RuO_4 510
Fermi liquid 39
Flux creep 446
–in $Pb_2Sr_2Y_{1-x}Ca_xCu_3O_8$ single crystals 442
–in $YBa_2Cu_3O_{7-x}$ single crystals 389
Flux depinning
–in $Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystal 412
Flux lattice elasticity
–in $La_{1.85}Sr_{0.15}CuO_4$ single crystal 420

Flux lattice melting
–in $YBa_2Cu_3O_y$ single crystals 409
Flux lattice 164, 412
Flux pinning 12, 164, 446
–in $Bi_2Sr_2CaCu_2O_{8+x}$ single crystal 431
–in $La_{1.85}Sr_{0.15}CuO_4$ single crystal 420
–in Nd123 crystals 396
Fluctuation effects 44

Gap energy 530
Gapless superconductivity 253
Ginzburg–Landau theory 21, 25

Hall effect 78, 138
–in κ-(BEDT-TTF)$_2$Cu(NCS)$_2$ 534
–in $YBa_2Cu_3O_7$ 381
–of $Bi_2Sr_2CuO_6$ single crystals 321
–of $La_{6-x}Sr_xBaLnCu_8O_{20}$ 325
–of $R_{1-x}Pr_xBa_2Cu_3O_7$ 333
Hamiltonian
–for $Y_{2-x}Ca_xBaNiO_5$ 118
Heat capacity
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystals 434
Heavy-fermion compounds 490, 498
High pressure synthesis 146
Hole Doping 193
Homologous series 146
Hubbard model 61, 66, 70, 74, 78, 86, 90, 111
Hund coupling 138, 142
Hybridization
–in $PrBa_2Cu_3O_{7-\delta}$ 340

I-V characteristics
–of $Bi_2Sr_2CaCu_2O_y$ single crystals 453
–of $Bi_2Sr_2CaCu_2O_8$ single crystals 461
IPES
–of $Bi_2Sr_2Ca_{1-x}Gd_xCu_2O_y$ 286
Impurity effect 94
Incommensurable magnetism 70
Inpurity effect 367
Intercalation 329
Interlayer hopping 54
Intrinsic Josephson junction
–of $Bi_2Sr_2CaCu_2O_y$ single crystals 453

Irreversibility line
–in $Pb_2Sr_2Y_{1-x}Ca_xCu_3O_8$ single crystals 442
Itinerant Heisenberg model 99

t–J model 35, 61, 94, 104
Josephson effect 238
–in $Bi_2Sr_2CaCu_2O_{8+x}$ single crystals 416
Josephson plasma mode
–in $Bi_2Sr_2CaCu_2O_8$ 381
Josephson plasma 1, 457
Junctions 218

Knight shift 526
–in $CuGeO_3$ 486
Kondo effect 490

Landau parameter 74
Logarithmic interaction 122
London penetration depth
–in $Tl_2Ba_2CaCu_2O_y$ 438
–in $YBa_2Cu_3O_y$ 438

Magnetic order
–in $La_{2-x}Ba_xCuO_4$ 290
Magnetic susceptibility
–of $Li(Ti_{1-x}V_x)_2O_4$ 523
–of $Sr_{14-x}Ba_xCu_{24}O_{41}$ 482
–of $Sr_{14-x}Ca_xCu_{24}O_{41}$ 482
–of $YSr_2Cu_2GaO_7$ 359
Magnetization 21, 90, 107, 446
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystals 434
–of $Pb_2Sr_2Y_{1-x}Ca_xCu_3O_8$ single crystals 442
–of $YBa_2Cu_3O_y$ single crystals 409
–of $YBa_2Cu_4O_8$ 464
Magnetoresistance 138
–of (La,Sr)MnO_3 134
–of $Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_2O_{8+\delta}$ single crystal 313
–of $Bi_2Sr_2CuO_6$ single crystals 321
–of $Nd_{2-x}Ce_xCuO_4$ thin film 317
–of $YBa_2Cu_3O_y$ crystals 302
Maximum entropy method 176, 340
Metal–insulator transition 49, 78, 122, 544
–in $(Sr_{0.4}Ca_{0.6})_{14}Cu_{24-x}Co_xO_{41-\delta}$ 475
–in $Bi_2Sr_2Ca_{1-x}Gd_xCu_2O_y$ 286
–of $La_{1.17-x}Sr_xVS_{3.17}$ 554
Misfit-layer compounds 554
Mixed state 25
Mott transition 111

NMR 378
–in $Bi_2Sr_2CaCu_2O_8$ single crystal 371
–in $CuGeO_3$ 486
–in $HgBa_2Ca_2Cu_3O_{8+\delta}$ 375
–of $(La_{1-y}Y_y)_{2-x}Ce_xCuO_4$ 355
–of $La_{2-x}Ba_xCuO_4$ 290
–of RT_2B_2C 526
–on $TlBa_2CaCu_2O_{7+\delta}$ 253
Nesting 378
Neutron diffraction
–of $Ca_2Sr_2Cu_3GaO_9$ 200
–of $La_{2-y-x}Bi_yBa_xCuO_4$ 298
Neutron inelastic scattering
–on $Y_{0.52}Pr_{0.48}Ba_2Cu_3O_7$ 333
Neutron powder diffraction
–of YBCO 176
Neutron scattering
–of $BaCo_{1-x}Ni_xS_2$ single crystal 550
–of $La_{2-x}Sr_xCuO_4$ 349
–of $YBa_2Cu_3O_{6+x}$ 257
Non Fermi liquid 490
Nonlinear Meissner effect 438
Nuclear density
–of YBCO 176

Optical conductivity
–of $YBa_2(Cu_{1-z}Zn_z)_3O_{7-y}$ single crystal 264
Organic superconductor 534, 538
Oxygen stoichiometry
–in $Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystals 434
–in $Ln_{1-x}Pr_xBa_2Cu_3O_{7-y}$ 336
–of La_2CuO_4 thin films 160
–of YBCO 197

Pair breaking
–in $Li(Ti_{1-x}V_x)_2O_4$ 523
Pairing mechanism 232
Peak effect
–in Nd123 crystals 396
–in Sr_2RuO_4 519
Phase diagram
–of κ-(BEDT-TTF)$_2$Cu(NCS)$_2$ 534
–of $BaCo_{1-x}Ni_xS_2$ 550
–of $La_{1-x}Sr_xMnO_3$ 544
–of $Pr_{1-x}Ca_xMnO_3$ 544
–of $Pr_{1-x}Sr_xMnO_3$ 544
–of $Sr_2CuO_2(CO_3)_{1-x}(BO_3)_x$ 344
–of $U_{1-x}Th_xPd_2Al_3$ 490
–of UPd_2Al_3 498
–of $Y_{1-x}U_xPd_3$ 490
Phase locking 453
Phase separation 44, 264
Phonon anomaly
–in $YBa_2Cu_3O_{6+x}$ 257
Phonon mechanism 126
Phonon scattering 305, 309
Phonon 363
Photoemission spectroscopy
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 208
Plaquette singlet
–in CaV_4O_9 114
Power spectrum 431
Preparation
–of Y123 filaments 450
Pseudogap
–in $YBa_2Cu_3O_{7-\delta}$ 277

Quantum Monte Carlo 90
Quantum oscillations
–in Sr_2RuO_4 510
Quasi-two-dimensional system 122

RPA 99
RVB 54
Raman scattering
–in $Pb_2Sr_2Ho(Y)_{1-x}Ca_xCu_3O_{8+\delta}$ 363
–in Sr_2RuO_4 single crystals 516
Refrigerator cooled magnet 172
Renormalisation 218
Renormalization-group method 82

STM 214, 245
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 225
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 241
–of $Nd_1Ba_2Cu_3O_{7-y}$ 185
STS 245
Scaling theory 49
Shear mechanism 151
Single-crystal growth
–of $BaCo_{1-x}Ni_xS_2$ 550
Singlet pairing 277
Slave boson 74
Specific heat
–of $YBa_2Cu_3O_{7-\delta}$ 277
Spectral function 104
Spin 1 chain 118
Spin correlation 111
Spin fermion model 142
Spin fluctuations 99, 378
–in $Bi_2Sr_2CaCu_2O_8$ single crystal 371
–in $HgBa_2Ca_2Cu_3O_{8+\delta}$ 375
–in $La_{2-x}Sr_xCuO_4$ 349
Spin gap 82, 114, 257, 475
–in $YBa_2Cu_4O_8$ 367
–of CaV_4O_9 271
Spin magnetic susceptibility 70
Spin-Peierls transition
–in $CuGeO_3$ 486
Spin-lattice relaxation time T_1
–in RT_2B_2C 526
Structural phase transformation 290
Structure phase transition 344
Structure
–of $Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_{2.0}O_x$ 180
–of $Ca_2Sr_2Cu_3GaO_9$ 200
–of $La_{2-y-x}Bi_yBa_xCuO_4$ 298

–of $LaCuO_{2.5}$ 468
–of $Nd_1Ba_2Cu_3O_{7-y}$ 185
–of $Sr_2CaRu_2O_y$ 558
–of $Sr_2Cu_3O_5$ 468
–of $Sr_2CuO_2(CO_3)_{1-x}(BO_3)_x$ 344
–of $Sr_3Ru_2O_7$ 558
–of $SrCu_2O_3$ 468
Substitution effect
–in $Ln_{1-x}Pr_xBa_2Cu_3O_{7-y}$ 336
Substitution effects
–in $TlSr_2Ca_{1-x}Y_xCu_2O_y$ 189
Superconductor 126
Superconducting fluctuation 313
Superconducting gap symmetry 241
Superconducting gap 214
Surface resistance
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 457
Susceptibility
–of $YBa_2Cu_3O_{7-\delta}$ 277
Symmetry of order parameter 257
Synthesis
–of Bi(2223) superconducting wires 172
–of $Ca_2Sr_2Cu_3GaO_9$ 200
–of La_2CuO_4 thin films 160
–of $LaCuO_{2.5}$ 468
–of $Pb_{0.7}Cu_{0.3}Sr_2Y_{0.5}Cu_2O_y$ 204
–of $Sr_2CaRu_2O_y$ 558
–of $Sr_2Cu_3O_5$ 468
–of $Sr_3Ru_2O_7$ 558
–of $SrCu_2O_3$ 468

T_c-suppression 333
Thermal conductivity
–of $REBa_2Cu_3O_7$ 309
–of $YBa_2(Cu_{1-x}Zn_x)_3O_{7-\delta}$ 305
Thermal fluctuation 317
Thermoelectric power 298
–of $La_{6-x}Sr_xBaLnCu_8O_{20}$ 325
Torque magnetometry
–of $YBa_2Cu_3O_7$ single crystal 260
Transport properties
–of $La_{1.17-x}Sr_xVS_{3.17}$ 554
Tunneling junction
–in $Ba_{1-x}K_xBiO_3$ thin films 530
Tunneling spectoscopy 238
Tunneling 218, 245

Ultrasound velocity
–in $La_{1.85}Sr_{0.15}CuO_4$ single crystal 420
Untwinned
–$YBa_2Cu_3O_7$ single crystal 260
Upper critical field H_{c2} 25
–of Sr_2RuO_4 single crystal 519

van-Hove singularity
–in Sr_2RuO_4 505
Vortex dynamics 12, 17, 401
Vortex glass 424
Vortex melting 401
Vortex state
–of $Bi_2Sr_2CaCu_2O_{8+\delta}$ 457
Vortex structure 428
Vortex glass 461

Weak coupling 416

XPS
–of $Bi_2Sr_2Ca_{1-x}Gd_xCu_2O_y$ 286
–of $Bi_2Sr_2CaCu_2O_y$ 294
–of $TlSr_2Ca_{1-x}Y_xCu_2O_y$ 189

ELSEVIER

Physica C 263 (1996) 575–576

PHYSICA C

Materials Index to Volume 263

$Ba_{1-x}K_xBiO_3$ thin films
–tunneling junction 530
$BaCo_{1-x}Ni_xS_2$ single crystal
–neutron scattering 550
$BaCo_{1-x}Ni_xS_2$
–phase diagram 550
–single-crystal growth 550
Bi(2223) superconducting wires
–synthesis 172
$Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_{2.0}O_x$
–crystal growth 180
–structure 180
$Bi_{2.1}Sr_{1.9}Ca_{1.0}Cu_2O_{8+\delta}$ single crystal
–magnetoresistance 313
$Bi_2Sr_2Ca_{1-x}Gd_xCu_2O_y$
–IPES 286
–metal–insulator transition 286
–XPS 286
$Bi_2Sr_2CaCu_2O_y$ single crystals
–*I-V* characteristics 453
–intrinsic Josephson junction 453
$Bi_2Sr_2CaCu_2O_y$
–XPS 294
$Bi_2Sr_2CaCu_2O_z$
–AGI intercalation 193
$Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystal
–*c*-axis resistivity 164
–flux depinning 412
$Bi_2Sr_2CaCu_2O_{8+\delta}$ single crystals
–heat capacity 434
–magnetization 434
–oxygen stoichiometry 434
$Bi_2Sr_2CaCu_2O_{8+\delta}$
–Fermi surface 208
–photoemission spectroscopy 208
–STM 225
–STM 241
–surface resistance 457
–vortex state 457
$Bi_2Sr_2CaCu_2O_{8+x}$ single crystal
–flux pinning 431
$Bi_2Sr_2CaCu_2O_{8+x}$ single crystals
–critical current density 416
–Josephson effect 416

$Bi_2Sr_2CaCu_2O_8$ single crystal
–NMR 371
–spin fluctuations 371
$Bi_2Sr_2CaCu_2O_8$ single crystals
–*I-V* characteristics 461
$Bi_2Sr_2CaCu_2O_8$
–anisotropy 381
–Josephson plasma mode 381
$Bi_2Sr_2CuO_6$ single crystals
–Hall effect 321
–magnetoresistance 321

$Ca_2Sr_2Cu_3GaO_9$
–neutron diffraction 200
–structure 200
–synthesis 200
CaV_4O_9
–plaquette singlet 114
–spin gap 271
$CuGeO_3$
–Knight shift 486
–NMR 486
–spin-Peierls transition 486

$HgBa_2Ca_2Cu_3O_{8+\delta}$
–NMR 375
–spin fluctuations 375

κ-(BEDT-TTF)$_2$Cu(NCS)$_2$
–electrical resistivity 538
–Hall effect 534
–phase diagram 534

(La,Sr)MnO_3
–magnetoresistance 134
$La_{1.17-x}Sr_xVS_{3.17}$
–metal–insulator transition 554
–transport properties 554
$La_{1.85}Sr_{0.15}CuO_4$ single crystal
–flux lattice elasticity 420
–flux pinning 420
–ultrasound velocity 420
$La_{1.85}Sr_{0.15}CuO_4$ single crystals
–energy gap 249

$La_{1-x}Sr_xMnO_3$
–phase diagram 544
La_2CuO_4 thin films
–oxygen stoichiometry 160
–synthesis 160
$La_{2-x}Ba_xCuO_4$
–magnetic order 290
–NMR 290
$La_{2-x}Sr_xCuO_4$
–detwinning 282
–electrical resistivity 282
–electronic specific heat 225
–neutron scattering 349
–spin fluctuations 349
$(La_{1-y}Y_y)_{2-x}Ce_xCuO_4$
–NMR 355
$La_{2-y-x}Bi_yBa_xCuO_4$
–neutron diffraction 298
–structure 298
$La_{6-x}Sr_xBaLnCu_8O_{20}$
–electrical resistivity 325
–Hall effect 325
–thermoelectric power 325
$LaCuO_{2.5}$
–structure 468
–synthesis 468
$Li(Ti_{1-x}V_x)_2O_4$
–electrical resistivity 523
–magnetic susceptibility 523
–pair breaking 523
$Ln_{1-x}Pr_xBa_2Cu_3O_{7-y}$
–oxygen stoichiometry 336
–substitution effect 336

Mn oxides
–effective Hamiltonian 130
–electronic structure 130

Nd123 crystals
–critical current density 396
–flux pinning 396
–peak effect 396

$Nd_1Ba_2Cu_3O_{7-y}$
–STM 185
–structure 185
$Nd_{2-x}Ce_xCuO_4$ thin film
–magnetoresistance 317

$Pb_{0.7}Cu_{0.3}Sr_2Y_{0.5}Cu_2O_y$
–electrical resistivity 204
–synthesis 204
$Pb_2Sr_2Ho(Y)_{1-x}Ca_xCu_3O_{8+\delta}$
–Raman scattering 363
$Pb_2Sr_2Y_{1-x}Ca_xCu_3O_8$ single crystals
–flux creep 442
–irreversibility line 442
–magnetization 442
$Pr_{1-x}Ca_xMnO_3$
–phase diagram 544
$Pr_{1-x}Sr_xMnO_3$
–phase diagram 544
$PrBa_2Cu_3O_{7-\delta}$
–hybridization 340

$R_{1-x}Pr_xBa_2Cu_3O_7$
–Hall effect 333
$REBa_2Cu_3O_7$
–thermal conductivity 309
RT_2B_2C
–NMR 526
–spin-lattice relaxation time T_1 526

$Sr_{14-x}Ba_xCu_{24}O_{41}$
–electrical resistivity 482
–magnetic susceptibility 482
$Sr_{14-x}Ca_xCu_{24}O_{41}$
–electrical resistivity 482
–magnetic susceptibility 482
$(Sr_{0.4}Ca_{0.6})_{14}Cu_{24-x}Co_xO_{41-\delta}$
–electrical resistivity 475
–metal–insulator transition 475
$Sr_2CaRu_2O_y$
–structure 558
–synthesis 558

$Sr_2Cu_3O_5$
–structure 468
–synthesis 468
$Sr_2CuO_2(CO_3)_{1-x}(BO_3)_x$
–phase diagram 344
–structure 344
Sr_2RuO_4 single crystal
–upper critical field H_{c2} 519
Sr_2RuO_4 single crystals
–Raman scattering 516
Sr_2RuO_4
–anisotropy 519
–electronic structure 505
–Fermi surface 505
–Fermi surface 510
–peak effect 519
–quantum oscillations 510
–van-Hove singularity 505
$Sr_3Ru_2O_7$
–structure 558
–synthesis 558
$SrCu_2O_3$
–structure 468
–synthesis 468

$Tl_2Ba_2CaCu_2O_y$
–London penetration depth 438
$TlBa_2CaCu_2O_{7+\delta}$
–NMR 253
$TlSr_2Ca_{1-x}Y_xCu_2O_y$
–charge transfer 189
–substitution effects 189
–XPS 189

$U_{1-x}Th_xPd_2Al_3$
–phase diagram 490
UPd_2Al_3
–phase diagram 498

Y123 filaments
–critical current density 450
–preparation 450
$Y_{2-x}Ca_xBaNiO_5$
–Hamiltonian 118

$Y_{0.52}Pr_{0.48}Ba_2Cu_3O_7$
–neutron inelastic scattering 333
$Y_{1-x}U_xPd_3$
–phase diagram 490
YBCO
–crystal growth 197
–neutron powder diffraction 176
–nuclear density 176
–oxygen stoichiometry 197
$YBa_2(Cu_{1-x}Zn_x)_3O_{7-\delta}$
–thermal conductivity 305
$YBa_2(Cu_{1-z}Zn_z)_3O_{7-y}$ single crystal
–optical conductivity 264
$YBa_2Cu_3O_y$ crystals
–magnetoresistance 302
$YBa_2Cu_3O_y$ single crystals
–flux lattice melting 409
–magnetization 409
$YBa_2Cu_3O_y$
–London penetration depth 438
$YBa_2Cu_3O_{6+x}$
–neutron scattering 257
–phonon anomaly 257
$YBa_2Cu_3O_7$ single crystal
–torque magnetometry 260
–untwinned 260
$YBa_2Cu_3O_7$
–electrical resistivity 401
–Hall effect 381
$YBa_2Cu_3O_{7-\delta}$ film
–current-voltage characteristics 424
$YBa_2Cu_3O_{7-\delta}$
–pseudogap 277
–specific heat 277
–susceptibility 277
$YBa_2Cu_3O_{7-x}$ single crystals
–columnar defects 389
–flux creep 389
$YBa_2Cu_4O_8$
–electrical resistivity 464
–magnetization 464
–spin gap 367
$YSr_2Cu_2GaO_7$
–antiferromagnetic order 359
–magnetic susceptibility 359